

SNRs in the Galactic Center region observed with *ASCA*

M. Sakano^{1,3}, J. Yokogawa¹, H. Murakami¹, K. Koyama^{1,5}, Y. Maeda^{2,4} and The *ASCA* Galactic Plane/Center Survey team

¹ Department of Physics, Kyoto University, Sakyo, Kyoto 606-8502, Japan

² Department of Astronomy and Astrophysics, Pennsylvania State University, University Park PA 16802-6305 U.S.A.

³ Research Fellow of the Japan Society for the Promotion of Science

⁴ Postdoctoral Fellow for Research Abroad of JSPS

⁵ CREST: Japan Science and Technology Corporation (JST)

Abstract. We report the *ASCA* results of the supernova remnants (SNRs) and their candidates in the Galactic Center region. We found apparent X-ray emission from G359.1–0.5 and G0.9+0.1, and made marginal detection for G359.1+0.9, but found no significant X-ray from the other cataloged SNRs: G359.0–0.9, Sgr A East (G0.0+0.0), G0.3+0.0, Sgr D SNR (G1.0–0.1) (Green 1998). The emission from G359.1–0.5 is found to be thermal with multi temperature structures whereas that from G0.9+0.1 is quite hard, probably non-thermal. We discovered two new candidates of SNRs: G0.0–1.3 (AX J1751–29.6) and G0.56–0.01 (AX J1747.0–2828). The former, G0.0–1.3, shows the extended emission with a thin thermal plasma. The latter, G0.56–0.01, shows quite strong 6.7-keV line with the equivalent width of 2 keV, which resembles that of the Galactic Center plasma. We discuss the nature of those SNRs, relating with the origin of the Galactic Center hot plasma.

1. Introduction

The hard X-ray structure of the Galactic Center region is quite unusual as well as those of the other wavelengths. In particular, thin thermal hot plasma prevails all over the region (Koyama et al. 1989; Yamauchi et al. 1990; Koyama et al. 1996; Maeda 1998). It has several surprising natures. First, the temperature is quite high, about 10 keV. Second, the total energy is also high, about 10^{54} erg. Third, the plasma wide-spreads in the region bigger than $1^\circ \times 1^\circ$, which corresponds to about $150\text{pc} \times 150\text{pc}$ for the distance of the Galactic Center. Fourth, the spectrum of the plasma is surprisingly uniform from field to field except for the slight change of the surface brightness. The origin of the plasma is still an enigma.

Supernova remnants (SNRs) are one of the possible candidates for the origin of the hot plasma (e.g., Koyama et al. 1986), as well as the other candidates; the past activity of the central massive black hole (e.g., Koyama et al. 1996), unresolved many cataclysmic variables (e.g., Mukai & Shiokawa 1993), and the global mag-

netic activity which heats the interstellar matter (e.g., Yokoyama et al. 1998).

If the SNRs explain all the hot plasma, first, the total energy ($\sim 10^{54}$ erg) requires about 10^3 supernovae in the narrow region with the scale of a few hundred pc in the last 5×10^4 years, which is the age of the plasma (Koyama et al. 1996). It would lead to the much higher supernova rate than the common rate of a few supernovae in a century in all the Galaxy. However, the possible starburst activity in the Galactic Center region has been often discussed. In particular, recent *COMPTEL* observations of ^{26}Al 1.8 MeV line suggest that 10^5 supernovae occurred in the last 4×10^6 years (Chen et al. 1995; Hartmann 1995), which is consistent with the required supernova rate.

Second, the supernova origin hypothesis requires the mean temperature of about 10 keV. It is much higher than the usual temperature of SNRs ($kT \leq$ a few keV). However, if SNRs densely exist in the region, the temperature may heat up to about 10 keV by the mutual interaction of the shocks. Thus, SNRs are still one of the candidates for the origin of the Galactic Center plasma.

In this paper we concentrate on the X-ray natures of the SNRs in the Galactic Center region from the observational point of view. The *ASCA* capability of imaging spectroscopy with 0.5–10 keV (Tanaka et al. 1994) is quite suitable for the study. In fact, owing to the hard X-ray sensitivity of *ASCA*, particularly above 2 keV, we can possibly detect new SNRs which have been hidden by the heavy interstellar absorption, hence obtain some information whether the SNRs are valid for the origin of the Galactic Center plasma.

Using all the available *ASCA* pointing observations in this region ($-1^\circ < l < 1^\circ$, $-1^\circ < b < 1^\circ$) and a part of the data of the ongoing *ASCA* Galactic Center survey, we studied the X-ray emission from the known SNRs (Green 1998), and searched new SNRs. We assume the distance to the Galactic Center to be 8.5 kpc.

2. Results on the cataloged SNRs

We searched X-ray emission from the cataloged SNRs (Green 1998), and then investigated each nature where significant X-ray emission is detected.

2.1. G359.1–0.5

Fig. 1 (upper panel) shows the *ASCA* GIS image of G359.1–0.5 with 1.6–2.1 keV band. We detected center-filled X-rays from this source, whereas the radio image shows a clear shell-like structure (e.g., Uchida et al. 1992).

The spectrum is found to have emission lines from highly ionized ion (Fig. 1 (lower panel)), hence the X-rays come from thin thermal plasma. The most distinct two lines are $K\alpha$ lines from helium-like silicon and hydrogen-like sulfur. They imply that the plasma has multi-temperature structure. The absorption column density is estimated at $N_H \sim 8 \times 10^{22} \text{ H cm}^{-2}$, suggesting that G359.1–0.5 is located at near the Galactic Center, which is consistent with the radio observations (Uchida et al. 1992). This column density is larger by a factor of 2 or 3 than that with the *ROSAT* measurement (Egger & Sun 1998). However, we are still confident of our estimation because the *ASCA* energy band is the most suitable for measuring such heavy absorption. Details of the analysis are given in Yokogawa et al. (1999).

2.2. G0.9+0.1

Fig. 2 (upper panel) shows the GIS image of G0.9+0.1 with 3–10 keV band. We detected significant X-ray emission from this source in this hard energy band, whereas no significant X-ray is detected in the softer energy band.

The X-ray emitting region is compact and not resolved with *ASCA* GIS. The radio size of the source is $2'$ in diameter (Helfand & Becker 1987), hence it is consistent with the X-ray image.

The spectrum is found to be hard; $kT > 2 \text{ keV}$ (the best-fit of 40 keV) in thermal model or $\Gamma \sim 1.5$ in power-law model, with heavy absorption of $N_H \sim 10^{23} \text{ H cm}^{-2}$, although the statistics are not good. This hardness may imply the emission to be non-thermal origin. The flux is $2 \times 10^{-12} \text{ erg s}^{-1} \text{ cm}^{-2}$ in 2–10 keV band. It is consistent with the upper limit by *Einstein* (Helfand & Becker 1987), but significantly lower than the *SAX* result (Mereghetti et al. 1999).

2.3. G359.1+0.9

Fig. 3 (upper panel) shows the GIS image around G359.1+0.9. X-ray emission from the position corresponding to G359.1+0.9 is detected with the significance of 9.8σ in 0.7–3 keV band (AX J1739.6–2911 in Fig. 3), whereas the significance in 3–10 keV band is 2.1σ .

Faint extended emission around AX J1739.6–2911 also can be seen in the 0.7–3 keV band image. We made the radial profile with the center at the peak of AX J1739.6–2911 and fitted it with the model of the point spread function (PSF) and the background (NXB+CXB(cosmic X-ray background)) where the normalizations of the PSF and the background were allowed to be free. The model is found to be rejected with the confidence of 97.4%, *i.e.*, the slightly extended emission with the radius of $4' \sim 5'$ was marginally detected with the significance of 2.2σ (Fig. 3 lower panel). The size is consistent with the radius of the radio shell, $r \sim 5'$.

The spectrum of the central region of AX J1739.6–2911 is found to be well fitted with the thin thermal plasma model with $kT \sim 0.7 \text{ keV}$ ($\chi^2/\text{d.o.f.} = 2.67/5$) and $N_H \sim 0$, although the statistics are not good. The flux is $3 \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$ in total energy band, converted to the luminosity of $2 \times 10^{33} \text{ erg s}^{-1}$ if we assume the distance to be 8.5 kpc. Note that the soft spectrum with no absorption may imply that this source is not located in the Galactic Center region, but is a foreground source, hence the above estimated luminosity may be reduced by several factors or more.

The extended emission with positional coincidence with the radio structure strongly supports that it is an X-ray counter part of the SNR, if the detection of the extended emission is true. The soft spectrum also supports it. The luminosity may be rather low for an SNR. However, *ASCA* Plane survey has failed to detect many radio SNRs, suggesting their quite dim X-ray luminosities (Yamauchi & *ASCA* Galactic Plane Survey team 1999). Therefore, the low X-ray luminosity of this source may be acceptable for an SNR. Future observations with high sensitivity will be encouraged.

2.4. The other cataloged SNRs

For the other cataloged SNRs (G359.0–0.9, Sgr A East (G0.0+0.0), G0.3+0.0, Sgr D SNR (G1.0–0.1)), we detected no significant X-ray emission associated with the SNRs. It is possibly because no strong X-ray is actually emitted. However we should be conservative for no X-ray emission from those sources because the detected positions of those SNRs are heavily contaminated from nearby strong X-ray sources; SLX 1744–299/300 near G359.0–0.9, Sgr A and AX J1745.6–2901 near Sgr A East, 1E 1743.1–2843 near G0.3+0.0, and GX3+1 near G1.0–0.1.

3. Results on the new candidates of SNRs

We discovered two new candidates of SNRs. In this section, we report the preliminary results on them.

3.1. G0.0–1.3 (AX J1751–29.6)

Fig. 4 (upper panel) shows the GIS image of G0.0–1.3 (AX J1751–29.6) in 0.7–3 keV band. We discovered the clearly extended emission with the scale of about $40' \times 10'$.

The spectrum is found to have the emission lines from highly ionized ions, hence is the thermal origin (Fig. 4 lower panel). In fact, the spectrum is well fitted with a thin thermal plasma model with $kT = 0.5 \pm 0.08$ keV and $N_H = (1.3 \pm 0.2) 10^{22}$ H cm $^{-2}$. The X-ray flux is $\sim 10^{-11}$ erg s $^{-1}$ cm $^{-2}$ in 0.5–3 keV band.

The column density suggests that this source may be in front of the Galactic Center region according to Sakano et al. (1999), and be located at the distance of about 4 kpc if we assume the mean interstellar density of 1 H cm $^{-3}$. Then, the obtained flux is converted to the luminosity of 3×10^{35} erg s $^{-1}$ under the assumption of the distance of 4 kpc. Even in the case of the quite small distance of 1 kpc, the luminosity is larger than 10^{34} erg s $^{-1}$.

We now consider the classification of the source. The clearly extended emission of the thin thermal plasma with $kT \sim 0.5$ keV implies that this source is an SNR or a star forming region. The luminosity is also within the typical range of SNRs but much higher than that of a star forming region. Therefore, AX J1751–29.6 is a strong candidate for a new SNR.

3.2. G0.56–0.01 (AX J1747.0–2828)

Fig. 5 (upper panel) shows the GIS image of G0.56–0.01 (AX J1747.0–2828) in 6–7 keV band, where this source can be seen the most significantly. The X-ray emitting region is compact and not resolved with GIS.

The background subtracted spectrum is given in Fig. 5 (lower panel). We accumulated the source X-ray photons from the $3'$ -radius circular region around AX J1747.0–2828, and the background photons, from an elliptical region with the major axis parallel to the Galactic Plane, excluding the $3'$ -radius circular regions around AX J1747.0–2828 itself and Sgr B2 (see Murakami et al. (1999)).

The spectrum is found to be characterized with a quite strong line at between 6–7 keV, which is also implied from the X-ray image. We tried to fit the spectrum with the thermal bremsstrahlung and a Gaussian line. Then we found the equivalent width of the line to be quite large, ~ 2 keV, and the center energy of the Gaussian to be 6.63 ± 0.06 keV, being consistent with $K\alpha$ line from helium-like iron. Hence, the spectrum is definitely a thin thermal origin with high temperature of several keV or higher.

We then fitted the spectrum with a thin thermal plasma model. The model well represents the total spectral shape. The best-fit temperature is $kT = 6.0^{+1.9}_{-1.5}$ keV, the abundance, $Z > 2$ solar, and the hydrogen column density, $N_H = (6.1^{+1.6}_{-1.1}) 10^{22}$ H cm $^{-2}$. The flux is 1.6×10^{-12} erg s $^{-1}$ cm $^{-2}$ in 0.7–10 keV band.

The large column density suggests this source to be located at near the Galactic Center. Thus, the absorption corrected X-ray luminosity is estimated at $\sim 3.6 \times 10^{34}$ erg s $^{-1}$ under the assumption of the distance of 8.5 kpc.

This high temperature and the overabundance suggest AX J1747.0–2828 to be a possible new candidate of a young SNR. The luminosity is also within the range of that of the typical SNR. Although the supernova remnant is the most probable source, the other possibility, for example, a cataclysmic variable, still cannot be excluded (e.g., Terada et al. 1999). In any case, it is a new class object in the Galactic Center region, which is a good candidate for the origin of the Galactic Center plasma.

4. Discussion

SNRs are one of the possible candidates for the origin of the Galactic Center hot plasma.

With *ASCA*, we completely surveyed the region of $|l| < 1^\circ$ and $|b| < 0.3^\circ$, where a large portion of SNRs in this region are expected to exist, and partially surveyed some areas with larger galactic latitude. The number of the detected SNRs with *ASCA* is two or three in 7 cataloged SNRs, and two for the new candidates in the Galactic Center region. On the other hand, the required number to be responsible for the Galactic Center plasma is about 10^3 in the region of $|l| \leq 1^\circ$ and $|b| \leq 0.5^\circ$. Therefore, the number of the detected SNRs is quite insufficient.

From recent radio molecular line observations, Hasegawa et al. (1998) found over 300 shell-like structures, possibly SNRs, in the region of $|l| \leq 0.5^\circ$, which correspond to about 20 shell-like structures along any line of sight in the region. Therefore, the number of SNRs may be possibly too much to resolve with *ASCA*. Future observations with higher angular resolution would be required to solve this problem.

The spectra of four of the detected five SNRs (or candidates) are relatively soft (G359.1–0.5, G359.1+0.9, and G0.0–1.3 (AX J1751–29.6)) or have no line-like feature (G0.9+0.1). Thus they cannot explain the spectrum of the Galactic Center plasma. On the other hand, a new SNR candidate G0.56–0.01 (AX J1747.0–2828) is notable for the strong iron line similar to that of the Galactic Center plasma. Such class of objects may be a key to understand the origin of the plasma. Search for such objects will be strongly encouraged.

Acknowledgements. The authors express their thanks to all the members of the *ASCA* team. We are grateful to Drs. J. P. Hughes and P. Slane for their valuable comments. MS and YM acknowledge the support from Research Fellowships of the Japan Society for the Promotion of Science.

References

Chen, W., Gehrels, N., Diehl, R., 1995, ApJ 440, L57

- Egger, R., Sun, X., 1998, In: “The Local Bubble and Beyond”, Breitschwerdt D., Freyberg M.J., Trümper J. (eds.), Springer, IAU Colloq. 166, 417
- Green, D.A., 1998, In: “A Catalog of Galactic Supernova Remnants”, <http://www.mrao.cam.ac.uk/surveys/snrs/>
- Hartmann, D.H., 1995, *ApJ* 447, 646
- Hasegawa, T., Oka, T., Sato, F., In: The Physics and Chemistry of the Interstellar Medium (Abstract Book), Aachen V.O.S.-V.
- Helfand, D.J., Becker, R.H., 1987, *ApJ* 314, 203
- Koyama, K., Awaki, H., Kunieda, H., et al., 1989, *Nature* 339, 603
- Koyama, K., Ikeuchi, S., Tomisaka, K., 1986, *PASJ* 38, 503
- Koyama, K., Maeda, Y., Sonobe, T., et al., 1996, *PASJ* 48, 249
- Maeda, Y., 1998, Ph.D thesis, Kyoto University
- Mereghetti, S., Sidoli, L., Israel, G.L., 1998, *A&A* 331, L77
- Mukai, K., Shiokawa, K., 1993, *ApJ* 418, 863
- Murakami, H., Koyama, K., Sakano, M., Maeda, Y., 1999, this volume
- Sakano, M., Koyama, K., Nishiuchi, M., et al., 1999, *Advances in Space Research*, in press
- Tanaka, Y., Inoue, H., Holt, S.S., 1994, *PASJ* 46, L37
- Terada, Y., Kaneda, H., Makishima, K., et al., 1999, preprint
- Uchida, K.I., Morris, M., Yusef-Zadeh, F., 1992, *AJ* 104, 1533
- Yamauchi, S., *ASCA* Galactic Plane Survey team, 1999, this volume
- Yamauchi, S., Kawada, M., Koyama, K., et al., 1990, *ApJ* 365, 532
- Yokogawa, J., Sakano, M., Koyama, K., Yamauchi, S., 1999, *Advances in Space Research*, in press
- Yokoyama, T., Tanuma, S., Kudoh, T., Shibata, K., 1998, In: The central regions of the Galaxy and galaxies, Sofue Y. (ed.), Kluwer Academic Publishers, IAU Symp. 184, 355




Fig. 1. (Upper) GIS contour map with 1.6–2.1 keV band superposed on a schematic diagram of the radio structures; the radio shell of G359.1–0.5 and the radio non-thermal filament, the Snake. Contour level is linearly spaced and is saturated for A1742–294 and 1E 1740.7–2942. The accumulated regions for the spectra are also noted. (Lower) The background-subtracted GIS spectrum. We also show the best-fit model where we fit the spectrum with the model of the thermal bremsstrahlung and two narrow Gaussians with interstellar absorption. These two figures are adopted from Yokogawa et al. (1999).




Fig. 2. (Upper) GIS2+GIS3 contour map of G0.9+0.1 with 3–10 keV band. The image is smoothed with a Gaussian filter of $\sigma = 0.75'$, and corrected for exposure, vignetting and the GIS grid structure after subtraction of non-X-ray background (NXB). Contour level is linearly spaced. The coordinate is in galactic (l_{II}, b_{II}) , and the north is up. (Lower) The same as Fig. 1 lower panel, but of G0.9+0.1. The fitting model is an absorbed power-law.




Fig. 3. (Upper) The same as Fig. 2 upper panel, but of G359.1+0.9 with 0.7–3.0 keV band. The positions of the detected X-ray sources (AX J1738.4–2903, AX J1739.3–2924, AX J1739.6–2911, AX J1740.3–2904), the radio shell of G359.1+0.9, and the radio pulsar PSR B1736–29 are indicated. (Lower) The radial profile with the center at the peak of AX J1739.6–2911. The profile is fitted with the point spread function (see text). The best-fit model is given with the dashed line, whereas the dotted line shows only the background component in the best-fit model.




Fig. 4. (Upper) The same as Fig. 2 upper panel, but of G0.0–1.3 (AX J1751–29.6) with 0.7–3.0 keV band. (Lower) The same as Fig. 1 lower panel, but of G0.0–1.3. For the background spectrum, we accumulated the photons from the elliptical region surrounding the source, excluding the source region, in the same GIS field of view. The fitting model is the thin thermal plasma model with absorption.




Fig. 5. (Upper) The same as Fig. 2 upper panel, but of G0.56–0.01 (AX J1747.0–2828) with 6.0–7.0 keV band, which is dominated by iron $K\alpha$ line. Note that the image was corrected only for exposure, NXB was not subtracted, and the region around a bright source 1E 1743.1–2843 was excluded before the smoothing in order to reduce the contamination from 1E 1743.1–2843. The positions of the X-ray reflection nebula Sgr B2 cloud (e.g., Murakami et al. 1999) and 1E 1743.1–2843 are also indicated. (Lower) The same as Fig. 1 lower panel, but of G0.56–0.01. The fitting model is the thin thermal plasma model and 6.4-keV narrow line, both with interstellar absorption.